

# Rapidity Dependence of High $p_T$ Suppression in Au+Au and d+Au Collisions at $\sqrt{s_{NN}}=200$ GeV.

I. G. Bearden<sup>7</sup>, D. Beavis<sup>1</sup>, C. Besliu<sup>10</sup>, Y. Blyakhman<sup>6</sup>, B. Budick<sup>6</sup>, H. Bøggild<sup>7</sup>, C. Chasman<sup>1</sup>, C. H. Christensen<sup>7</sup>, P. Christiansen<sup>7</sup>, J. Cibor<sup>3</sup>, R. Debbé<sup>1</sup>, E. Enger<sup>12</sup>, J. J. Gaardhøje<sup>7</sup>, M. Germinario<sup>7</sup>, K. Hagel<sup>8</sup>, O. Hansen<sup>7</sup>, A. Holm<sup>7</sup>, A. K. Holme<sup>12</sup>, H. Ito<sup>11</sup>, A. Jipa<sup>10</sup>, F. Jundt<sup>2</sup>, J. I. Jørdre<sup>9</sup>, C. E. Jørgensen<sup>7</sup>, R. Karabowicz<sup>4</sup>, T. Keutgen<sup>8</sup>, E. J. Kim<sup>1</sup>, T. Kozik<sup>4</sup>, T. M. Larsen<sup>12</sup>, J. H. Lee<sup>1</sup>, Y. K. Lee<sup>5</sup>, G. Løvholden<sup>12</sup>, Z. Majka<sup>4</sup>, A. Makeev<sup>8</sup>, B. McBreen<sup>1</sup>, M. Mikelsen<sup>12</sup>, M. Murray<sup>8</sup>, J. Natowitz<sup>8</sup>, B. S. Nielsen<sup>7</sup>, J. Norris<sup>11</sup>, K. Olchanski<sup>1</sup>, J. Olness<sup>1</sup>, D. Ouerdane<sup>7</sup>, R. Płaneta<sup>4</sup>, F. Rami<sup>2</sup>, C. Ristea<sup>10</sup>, D. Röhrich<sup>9</sup>, B. H. Samset<sup>12</sup>, D. Sandberg<sup>7</sup>, S. J. Sanders<sup>11</sup>, R. A. Scheetz<sup>1</sup>, P. Staszal<sup>7</sup>, T. S. Tveter<sup>12</sup>, F. Videbæk<sup>1</sup>, R. Wada<sup>8</sup>, A. Wieloch<sup>4</sup>, Z. Yin<sup>9</sup>, I. S. Zgura<sup>10</sup>

The BRAHMS Collaboration

<sup>1</sup> Brookhaven National Laboratory, Upton, New York 11973, USA

<sup>2</sup> Institut de Recherches Subatomiques and Université Louis Pasteur, Strasbourg, France

<sup>3</sup> Institute of Nuclear Physics, Krakow, Poland

<sup>4</sup> Smoluchowski Inst. of Physics, Jagiellonian University, Krakow, Poland

<sup>5</sup> Johns Hopkins University, Baltimore 21218, USA

<sup>6</sup> New York University, New York 10003, USA

<sup>7</sup> Niels Bohr Institute, Blegdamsvej 17, University of Copenhagen, Copenhagen 2100, Denmark

<sup>8</sup> Texas A&M University, College Station, Texas, 17843, USA

<sup>9</sup> University of Bergen, Department of Physics, Bergen, Norway

<sup>10</sup> University of Bucharest, Romania

<sup>11</sup> University of Kansas, Lawrence, Kansas 66049, USA

<sup>12</sup> University of Oslo, Department of Physics, Oslo, Norway

(Dated: Version: High-pt30.tex, June 25, 2003)

We present spectra of charged hadrons from Au+Au and d+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV measured with the BRAHMS experiment. The spectra are compared to spectra from nucleon-nucleon collisions at the same energy scaled by the number of binary collisions. The resulting ratios (nuclear modification factors) for central Au+Au collisions at  $\eta = 0$  and  $\eta \approx 2$  evidence a strong suppression in the high  $p_T$  region ( $>2$  GeV/c). In contrast the d+Au nuclear modification factor (at  $\eta = 0$ ) exhibits an enhancement of the high  $p_T$  yields. These measurements indicate a high energy loss of the high  $p_T$  particles in a medium created in the central Au+Au collisions. The lack of suppression in d+Au collisions rules out initial state effects as explanation for the suppression in the central Au+Au collisions.

PACS numbers: 25.75 Dw.

Collisions between heavy nuclei (eg. Au) in the energy domain now accessible at the Relativistic Heavy Ion Collider (RHIC) are expected to lead to the formation of an extremely hot high density region exhibiting features characteristic of quark deconfinement, the quark-gluon plasma (QGP). The first experiments with Au+Au collisions at 100 AGeV + 100 AGeV suggest that very high energy densities ( $\epsilon > 5 \text{ GeV}/fm^3$ ) are achieved in the initial stages of such collisions. Furthermore, the reaction mechanism at RHIC has new features as compared to lower energies, evidencing a high degree of nuclear transparency (as may be deduced from the low rapidity density of net protons that is measured in a region around midrapidity ( $|y| < 1.5$ ) ([1]). This picture brings to mind the boost invariant scenario for such collisions first proposed by Bjorken (ref), in which the colliding nuclei suffer only a moderate relative rapidity loss and where the subsequent particle production can be primarily traced to quark-antiquark pair production from the breaking

of color strings between scattered partons. Furthermore, studies of the particle production (ref. multiplicity papers from 4 experiments) and the dynamics of the expanding hadronic cloud that subsequently forms, suggest that the system, at least in later stages of the collision, may be in thermal and chemical equilibrium. At midrapidity, an analysis of particle ratios, indicates that the baryochemical potential has been reduced to 29 MeV [?, [8]. This set of observations naturally leads to speculation about whether a high density deconfined partonic state is indeed formed in Au+Au collisions at RHIC.

In order to investigate the conditions prevailing early in the evolution of the system it has been proposed (ref. Bjorken, Wang+Gulyassy etc...) that the study of high momentum scattered particles may be a good probe of the conditions of the original medium. Such particles, which are associated with jet production from initial hard parton scatterings, are predicted to experience attenuation and energy loss as they tra-

verse a dense hadronic medium with a high density of color charges due to induced gluon radiation, resulting in a reduction of the high transverse momentum component of their spectra. This process is referred to as  $p_T$  suppression.

In this Letter, we report on measurements of spectra of charged hadrons from Au+Au collisions at  $\sqrt{s_{NN}} = 200$  AGeV at pseudorapidities  $\eta = -\ln(\tan(\theta/2)) = 0$  and  $\eta = 2.2$ , where  $\theta$  is the angle of emission relative to the beam direction. The spectra, which have been measured as a function of the collision centrality, are compared to reference data from elementary  $p + \bar{p}$  collisions at the same energy and rapidity, using a scaling to the estimated number of binary collisions as described below. We have also measured similar spectra (for minimum bias collisions) for the reaction  $d + Au$  at  $\sqrt{s_{NN}} = 200$  AGeV. For central (0 – 10%) Au+Au collisions we find a strong suppression of the high transverse momentum component ( $p_t > 2\text{GeV}/c$ ) of the spectra as compared to the scaled  $p + \bar{p}$  spectra. This suppression diminishes significantly as the collision centrality decreases. In contrast no apparent suppression is observed for the  $d + Au$  collisions.

BRAHMS consists of two magnetic spectrometers (the MidRapidity Spectrometer, MRS and the Forward Spectrometer, FS) that together can cover the angular range from 2.3 degrees to 90 degrees relative to the beam direction and thus measure hadrons and antihadrons with rapidities in the range  $0 < y < 4$ . Particles are deflected in room temperature dipole magnets. Their directions before and after entering the region of vertical magnetic field is determined by the use of Time Projection Chambers and Drift Chambers. From the bending angle we determine the momentum of the particles. We estimate that the momentum resolution is (typically)  $\Delta p/p =$  at  $p = xx$  GeV/c and ... at .... The experiment also has excellent particle identification (PID) capabilities. In this work, we concentrate on the spectra of charged hadrons ( $h^+$  and  $h^-$ ), for which this PID capability is not used. In addition to the spectrometers a set of global detectors were used for minimum bias trigger and event characterization.

In the Au+Au run the data acquisition was triggered by a signal corresponding to one or more neutrons in each of the ZDC counters. Simulation studies using GEANT show that this corresponds to approximately 97% of the Au+Au interaction cross section. An additional trigger based on the charged particle multiplicity in the midrapidity region, selected the most central events (0 –  $\approx 25\%$ ) in parts of the Au+Au run. In the  $d + Au$  run scintillator counters (INEL) were placed around the nominal intersection point (IP) at  $z = +2$ , and  $+3$  m (check). Signals from these counters were used as the minimum bias trigger, which selected approximately 90% of the  $d + Au$  cross section. Additional spectrometer trigger were added to enhance




FIG. 1: Spectra of charged hadrons for centrality cuts 0–10, 10–20, 20–40 and 40–60 percent of the total reaction cross section for Au+Au collisions at  $\sqrt{s_{nn}} = 200$  AGeV and for pseudorapidities around  $\eta = 0$  (left panel) and  $\eta = 2.2$  (right panel). The left panel also shows the spectrum for elementary  $p + p$ -bar collisions from UA1, scaled to  $\eta = 0$ . All displayed spectra have been scaled by the indicated factor for clarity.

the track sample. Further details of the experimental setup and operation can be found in [4, 5, 9].

Centrality selection of the Au+Au collisions is done using global multiplicity detectors positioned around the nominal intersection point (IP). The position of the actual intersection point is determined with a precision of  $\sigma < 3$  cm by the use of arrays of beam counters (BB) placed at about 2m away on either side of the collision zone. For the  $d + Au$  reaction study the vertex measurement by the INEL counters has a resolution of approximately 9 cm. By combining the information from the vertex determined by the INEL counters, the BB counters and the tracks in first TPC of the midrapidity spectrometer the true vertex distribution was obtained and used for (vertex dependent) event normalization. In addition, fiducial cuts in the spectrometer opening is applied to avoid systematic errors due to edge effects.

Figure 1 shows measured spectra for charged hadrons ( $h^+ + h^-$ )/2 at 90 degrees (right panel) and for negative hadrons ( $h^-$ ) at 12 degrees (left panel), corresponding to pseudorapidities ( $\eta = 0$  and 2.2). The width of the selected rapidity intervals are (...) and (...) respectively. The displayed spectra correspond to various centralities (0%-10%, 10%-20%, 20%-40%, 40%-60%). The spectra have been constructed from measurements with various fields settings of the magnets in the MRS and FS respectively. Spectra have been corrected for the finite acceptance of the spectrometers, for the tracking efficiency, and where applicable, for the triggering efficiency. Spectra have also been corrected for effects due to the momentum resolution (FIX details...). Also shown in the figure is




FIG. 2: Top row: Nuclear modification factors  $R_{AuAu}$  as a function of transverse momentum for Au+Au collisions at  $\eta = 0$  and  $\eta = 2.2$  for 0 – 10% most central collisions. Middle row: as top row, but for centralities 40 – 60%. Bottom row: ratio of the  $R_{AuAu}$  factors at the two rapidities for the most central and most peripheral collisions. The dotted and dashed lines show the expected value using number of participants scaling and number of binary collisions scaling, respectively. Error bars are statistical. The grey bands indicate the systematic errors.

the ‘reference’ spectrum for  $p + \bar{p}$  collisions measured by the UA1 experiment at CERN-SPS (ref...). This spectrum was measured in the range  $|\eta| < 2.5$ . To be able to compare it to our data, which are confined to narrower rapidity intervals, we have based used a momentum dependent correction factor estimated with the use of the HIJING code (ref...). This code, based on PYTHIA, has been tuned to reproduce p+p collisions. We have compared our corrected spectrum at midrapidity with the 100AGeV+100AGeV  $p + p$  distribution, recently measured by the STAR collaboration (ref...). We find excellent agreement. No similar comparison is available for the more forward rapidity. Consequently we use the model-scaled  $p + \bar{p}$  spectrum for the following comparisons, noting that HIJING predicts a difference between negatively and positively charged hadrons at forward rapidities.

A useful way to compare the momentum spectra from nucleus-nucleus collisions to those from p+p collisions, is to scale the normalized p+p spectrum by the number of binary collisions ( $N_{bin}$ ) corresponding to the centrality cuts applied to the nucleus-nucleus spectra and construct the ratio. This ratio is called the nuclear modification factor,  $R_{AA}$  (FIX: Formula). The used  $N_{bin}$  values have been obtained from the HIJING code. For the various centrality cuts in Au+Au they are. .... and for the d+Au reaction we

have used  $N_{bin} = 7$ .




FIG. 3: Ratio of  $R_{cp}$  distributions at  $\eta = 0$  and  $\eta = 2.2$  from the bottom row of figure 3.

Figure 2 (upper two rows) shows the ratios  $R_{AuAu}$ , as a function of transverse momentum for different centrality cuts for the Au+Au measurements at  $\eta = 0$  and 2.2, respectively. The salient features of the distributions is a rise from about 0.2 in the range  $p_t = 0.5 - 2\text{GeV}/c$  to a value in the range 0.5 to 0.9. This part of the spectrum is normally associated with soft collisions and should therefore scale with the number of participants, wherefore a scaling with the (larger)  $N_{bin}$  reduces  $R_{AuAu}$  in this momentum interval. (FIX) Beyond  $p_T \approx 2\text{GeV}/c$ ,  $R_{AuAu}$  is expected to be close to 1. In fact measurements of  $R_{AuAu}$  at CERN-SPS for  $\sqrt{s_{NN}} = 17\text{ GeV}$  collisions for neutral pions (ref: WA98), negative hadrons (ref: NA49) and charged pions (ref: CERES) show that  $R_{AA}$  is equal to 1 at  $p_T = 1.5\text{ GeV}/c$  and increases to about 1.5 at  $p_T = 3\text{GeV}/c$ . This is called the Cronin effect ref. Cronin etc.) and explained as being due to multiple scattering of nucleons in the initial stages of the collision. Above  $p_T > 2\text{GeV}/c$  the  $R_{AA}$  distributions shown in fig. 2 are systematically lower than 1 for the central collisions at, while they tends towards 1 for the peripheral collisions. Indeed, for the most central collisions at both rapidities,  $R_{AuAu}$  is only about 0.4 at  $p_T \approx 3\text{ GeV}/c$ . The high  $p_T$  component of the Au+Au spectra is therefore suppressed by a factor of 3-4 as compared to the SPS results at lower energies. The two lower panels in fig.2 show the ratio of the  $R_{AuAu}$  distributions for the most central collisions relative to the least central at the two rapidities. We call this the  $R_{cp}$  ratio. Both ratio distributions clearly show a significant suppression of the high  $p_t$  component for the central collisions.

In figure 3 we present the ratio of the  $R_{cp}$  distributions from  $\eta = 0.0$  and  $\eta = 2.2$ . This ratio has the advantage that it is to a large extent free from systematic errors arising from the normalization to the




FIG. 4: Nuclear modification factor measured for near minimum bias collisions of D+Au at  $\sqrt{s_{nn}} = 200$  AGeV. Error bars represent statistical errors. Systematic errors on the ratio are 20percent

spectra from elementary collisions. The figure shows that, within errors, the degree of high  $p_T$  suppression observed at  $\eta = 0$  and at  $\eta = 2.2$  is similar or perhaps even superior at the more forward rapidities.

For comparison to the results obtained for the Au+Au collisions we have investigated the d+Au reaction at the same energy and at  $\eta = 0$ ). In figure 4 we present the corresponding  $R_{dAu}$  distribution, analyzed in the same way as the Au+Au collisions, and using the same reference spectrum, appropriately scaled by  $N_{bin}$ . We have applied no centrality cuts, so the distribution reflects our minimum bias collision data, which correspond to  $\approx 90\%$  of the total reaction cross section. We have verified that the shape of the  $R_{dA}$  distribution is roughly similar for different centrality cuts. It is striking that the  $R_{dAu}$  factor shows no suppression of the high  $p_T$  component, rather it shows an enhancement, as has been found at lower energies.

From these measurements we conclude that central collisions between Au+Au nuclei exhibit a very significant suppression of the high transverse momentum component as compared to  $p + p$  collisions. This

suppression appears to be directly correlated with the size of the participant zone, as demonstrated by the fact that the much smaller participant zone resulting from the d+Au collisions shows no suppression and by the fact that more peripheral collisions between the Au nuclei show less suppression than central collisions. It is reasonable to surmise that the effect is related to medium effects tied to a large volume with high energy density. It has been proposed that gluon saturation effects in the colliding Au+Au nuclei, i.e. entrance channel effects, resulting from their high laboratory energy, might limit the phase space available for the production of high momentum particles in the collisions. Such an explanation appears improbable in view of the results for the d+Au measurements which utilize projectiles (of which one is larger) at the same energy.

In summary, the BRAHMS measurements demonstrate a significant suppression of the high  $p_T$  component of transverse momentum spectra for hadrons measured at two rapidities. The suppression is seen to diminish with decreasing collision centrality and is seen to be absent in collisions between a light mass nucleus and a gold nucleus. We conclude that the observed suppression is consistent with significant medium effects in the most violent collisions, i.e. those that have the largest participant volumes. The fact that the suppression persists at forward rapidities suggests that the volume which causes the suppression is extended also in the longitudinal direction. Whether the observed effect is tied to absorption or energy loss of high momentum particles at the partonic level (i.e. in the very initial stages of the collisions) or to some other mechanism is as yet unclear and warrants further systematic investigations, notably by studying the effect over the largest possible rapidity range and by carrying out experiments at lower beam energies.

This work was supported by the division of Nuclear Physics of the Office of Science of the U.S. DOE, the Danish Natural Science Research Council, the Research Council of Norway, the Polish State Com. for Scientific Research and the Romanian Ministry of Research.

- 
- [1] Quark Matter 2002, BRAHMS collaboration Nucl. Phys. A715 (2003) xxx, and P. L. Christiansen, Ph. D. thesis, Univ. Copenhagen, June 2003.
  - [2] Quark Matter 2002, BRAHMS collaboration Nucl. Phys. A715 (2003)715.
  - [3] I. G. Bearden *et al.*, BRAHMS Collaboration, Phys. Rev. Lett. **90**, 112305 (2003)
  - [4] I. G. Bearden *et al.*, BRAHMS Collaboration, Phys. Lett. B **523**, 227 (2001).
  - [5] I. G. Bearden *et al.*, BRAHMS Collaboration, Phys. Rev. Lett. **88**, 202301 (2002).
  - [6] B. B. Back *et al.*, PHOBOS Collaboration, Phys. Rev. Lett. **88**, 022302 (2002).
  - [7] J. D. Bjorken, Phys. Rev. D**27**, 140 (1983).
  - [8] P. Braun-Munzinger *et al.*, Phys. Lett. B **518**, 41 (2001).
  - [9] M. Adamczyk *et al.*, BRAHMS Collaboration, Nucl. Instr. and Meth., A499 (2003) 437.