Proc. 23rd Winter Workshop on Nuclear Dynamics (2007) 000–000

23rd Winter Workshop on Nuclear Dynamics Big Sky, Montana, USA February 11–18, 2007

# Results from pp at 62.4 and 200 GeV with the BRAHMS experiment

# F.Videbæk<sup>1</sup> for the BRAHMS experiment

<sup>1</sup> Brookhaven National Labloratory, NY 11973 Upton, New York, US

Abstract. Measurements of elementary pp collisions is an integral component to understand heavy ion collisions. Results for pp collisions at 200 and 62.4 GeV are presented. At both energies NLO pQCD describes pion production well. The measured pion transverse single spin asymmetries are very large at 62.4 GeV and are reasonably well described by models relying on pQCD at transverse momenta larger than 1 GeV/c

*Keywords:* pp spectra, RHIC, single spin asymmetry *PACS:* 13.85.Ni,13.85.Hd,13.88+e,12.38Qk,25.75.-q

## 1. Introduction

One of the goals of the relativistic heavy ion program is to study the properties of matter at high temperature and high density. The explorations at the Relativistic Heavy Ion Collider (RHIC) indicate that that at c.m. energies of 200 GeV per nucleon pair indeed such system is formed with novel properties as characterized through the particle production [1–4]. Part of these conclusion relies on comparison to elementary pp collisions, where such nuclear effects should not be present. The RHIC experiments have also explored Au+Au and Cu+Cu collisions at 62.4 GeV, an intermediate energy between that at the SPS (18 GeV) and that where the bulk of the RHIC heavy ion program is performed at 200 GeV. The 62.4 GeV was chosen to match the highest energy where data were obtained in the ISR experiments. Thus, to minimize the systematic uncertainty for the comparisons to the heavy ion data a brief run with (polarized) pp at 62.4 GeV was carried out to collect pp reference data by the BRAHMS, PHENIX and STAR experiments. Thus, at this point in time extensive set of pp reference data have been collected at these two energies.

In addition to the pp comparison data there is an extensive spin program at RHIC. It had long been envisioned that BRAHMS could contribute to the studies using the transversely polarized beams with studies of pions at large momenta at forward angles. Part of run-5 and run-6 was dedicated to such measurements. There is current interest in understanding the large transverse single spin asymmetries (SSA) observed under these conditions in the energy range from 20 to 200 GeV in the framework of different approaches to pQCD. Main theoretical efforts to account for the observed has been on the role of transverse momentum dependent (TMD) partonic effects in the structure of the initial transversely polarized nucleon[5] and the fragmentation process of a polarized quark into hadrons[6]. Higher twist effects ("twist-3") arising from quark-gluon correlation effects have also been considered as a possible origin of SSA[7,8]. To validate such an approach that is based of pQCD it is key to see how well pQCD describes inclusive data at high rapidities.

The contribution presents results of inclusive pp reactions at 200 and 62.4 GeV, and comparisons to pQCD at both energies. Transverse spin results from 62.4 and 200 GeV are also described and compared to model calculations.

#### 2. Inclusive pp measurements

The data used for this analysis were collected with the BRAHMS detector system. The BRAHMS detector consists of two movable magnetic spectrometers, the Forward Spectrometer (FS) that can be rotated from  $2.3^{\circ}$  to  $15^{\circ}$ , and the Mid-Rapidity Spectrometer (MRS) that can be rotated from  $34^{\circ}$  to  $90^{\circ}$  degrees relative to the beam line, and several global detectors for measuring multiplicities, luminosity, and determining the interaction vertex, and provides a start time (T0) for time-of-flight measurement.

The MRS is a single-dipole-magnet spectrometer with a solid angle of  $\approx 5msr$ and a magnetic bending power up to 1.2 Tm. Most of the data presented here were recorded at magnetic field settings of 0.4 and 0.6 Tm. The MRS contains two time projection chambers (TPCs), TPM1 and TPM2 sitting in field free regions in front and behind the dipole (D5). This assembly is followed by a two highly segmented scintillator time-of-flight walls, one (TOFW) at 4.51 m and a second (TFW2) at either 5.58 m (90° setting) or 6.13 m (other angle settings).

The forward Spectrometer consists of 4 dipole magnets D1, D2, D3 and D4 with a bending power of up to 9.2 Tm. The spectrometer has 5 tracking stations T1 through T5, and particle identifiing detectors H2, a segmented time-of-flight wall, and the Ring Imaging Detector (RICH)[9]. Additional details on the BRAHMS experimental setup can be found in ref. [10].

The minimum bias trigger used to normalize these measurements is defined with a set of Cherenkov radiators (CC) placed symmetrically with respect to the nominal interaction point and covering pseudo-rapidities that range in absolute value from 3.26 to 5.25. This trigger required that at least one hit is detected in both sides of the array.

The present analysis was done with charged particles that originated from collisions of polarized protons with interaction vertices in the range of  $\pm 40$  cm. For the 200 GeV data invariant cross sections were extracted in narrow ( $\Delta y = 0.1$ ) rapidity bins centered at y=2.95 and y = 3.3, respectively. Narrow rapidity bins



Fig. 1. Invariant cross section distributions for pion, kaons protons and antiprotons produced in p+p collisions at  $\sqrt{s} = 200$  GeV at rapidities y=2.95 (left panels) and y = 3.3 (right panels). In all panels, positive charged particles are shown with filled triangles and negative ones with open circles. The errors displayed in these plots are statistical.

are required to reduce the effects of rapidly changing cross sections in particular at higher  $p_{\rm T}$ . Each distribution is obtained from the merging of up to five magnetic field settings. The data are corrected for the spectrometer geometrical acceptance, multiple scattering, weak decays and absorption in the material along the path of the detected particles. Tracking and matching efficiencies for each of the 5 tracking stations in the spectrometer were calculated by constructing full tracks with only 4 stations and evaluating the efficiency in the 5<sup>th</sup> station. The overall efficiency is about 80-90% and is included in the extraction of the cross sections. Particles are identified by the RICH. The low momentum part of the proton spectra is measured using the RICH in veto mode.

Data for y=2.95 and y=3.3 are presented for pions, kaons and protons in Fig.1. The pions exhibit a power law behavior, and at high  $p_{\rm T}$  the ratio  $\pi^-/\pi^+$  is less than one indicating the increasing importance of valence quark. The  $\bar{p}/p$  ratio is rather small at the higher  $p_{\rm T}$  indicating that in the case of fragmentation the gluons cannot dominate the production.

The measured differential cross-sections are compared with NLO pQCD calculations [11] evaluated at equal factorization and renormalization scales,  $\mu \equiv \mu_F = \mu_R = p_T$ . These calculations use the CTEQ6 parton distribution functions [12]



Fig. 2. Comparison to NLO pQCD for pp at 200 geV

and a modified version of the "Kniehl-Kramer-Potter" (KKP) set of fragmentation functions (FFs) [13] referred to here as mKKP, as well as the "Kretzer" (K) set [14]. The KKP set includes functions that fragment into the sums  $\pi^+ + \pi^-$  and  $K^+ + K^-$ . Modifications were necessary to obtain functions producing the separate charges for both  $\pi$  and K. These modifications involve the following operations: to obtain the FFs producing  $\pi^+$ , the functions fragmenting favored light quarks  $u, \bar{d}$ into  $\pi^0$  were multiplied by (1+z) (e.g.  $D_u^{\pi^+} = (1+z)D_u^{\pi^0}$  with  $D_u^{\pi^0} = \frac{1}{2}D_u^{\pi^+ + \pi^-}$ ) where z is the fraction of the parton momentum carried by the hadron, and  $D_u^{\pi^+}$ the function fragmenting u quarks into positive pions. The functions fragmenting unfavored quarks  $\bar{u}, d$  into  $\pi^0$  are multiplied by 1 - z. The same operation is done for  $\pi^-$ , but this time the favored quarks are  $\bar{u}$  and d. The FFs of strange quarks and gluons are left unmodified. Similar modifications were applied to obtain FFs into  $K^+$  and  $K^-$ , but this time, the starting functions were the ones fragmenting  $u, \bar{u}, s, \bar{s}$  into the sum  $K^+ + K^-$ . Figure 2 shows that the agreement between the NLO calculations that include the mKKP FFs and the measured pion cross section is remarkable (within 20% above 1.5 GeV/c). Similar good agreement was obtained for neutral pions at y=0 [15] and at y=3.8 [16] at RHIC. The agreement between the calculated and the measured kaon cross-sections is equally good. The difference between the mKKP and Kretzer parametrizations is driven by higher contributions from gluons fragmenting into pions. This difference has been identified as an indication that the gg and gq processes dominate the interactions at mid-rapidity [15]. The present results indicate that such continues to be the case at high rapidity. The calculation that uses the Kretzer set underestimates the pion yields by a factor of  $\sim 2$  at all values of  $p_{\rm T}$  while for positive kaons, the agreement is good at low momentum but deteriorates at higher momenta. Additional details as well as



Fig. 3. Comparison to NLO pQCD for pp at 62 GeV

comparisons to p and  $\bar{p}$  spectra can be found in Ref.[17].

At 62.4 GeV where the beam rapidity is 4.2 the spectrometer at forward angle samples produced particle that carries a significant fraction of the available momenta (31.1 GeV/c). Thus particle production is influenced by the kinematic limit. Data for identified charged hadrons were collected at 2.3°,3°, 4°, and 6°. In Fig3 for rapidities 2.7 and 3.3. The cross sections changes rapidily with rapidity at high  $p_T$ where  $x_F$  values up to 0.5 are probed. The data are compared to NLO pQCD in the same figure. The calculation are done in the same manner as the 200 GeV using the KPP fragmentation function evalauted at  $\mu = p_T$  scale. The calculation describe the overall magnitude and shape quite well, though at the highest rapidity there is a tendency for the calculation to fall below the data at the highest  $p_T$ . This may be in agreement with the analysis[18] of 53 GeV  $\pi^0$  data from the ISR at a fixed angle of 4°, comparable to the present measurements (2.3° and 4°), but at larger  $x_F$ where NLO pQCD is considerable below the data and with increasing discrepancy with  $x_F$ . In contrast to the aforementioned paper we do though conclude that NLO pQCD gives a satisfactory description of the charged pion data at high rapidity.

#### 3. Transverse Single Spin Asymmetries

The SSA is defined as a "left-right" asymmetry of produced particles from the hadronic scattering of transversely polarized protons off unpolarized protons. Experimentally the asymmetry can be obtained by flipping the spins of polarized protons, and is customary defined as analyzing power  $A_N$ :

$$A_N = \frac{1}{\mathcal{P}} \frac{(N^+ - \mathcal{L}N^-)}{(N^+ + \mathcal{L}N^-)},$$
 (1)

where  $\mathcal{P}$  is polarization of the beam,  $\mathcal{L}$  is the spin dependent relative luminosity ( $\mathcal{L} = \mathcal{L}_+/\mathcal{L}_-$ ) and  $N^{+(-)}$  is the number of detected particles with beam spin vector



oriented up (down). The average polarization of the beam  $\mathcal{P}$  as determined from the

**Fig. 4.**  $A_N$  v.s  $x_F$  for  $\pi^+$  and  $\pi^-$  at 62 GeV for positive and negative  $x_F$ . Solid symbols are for  $\pi^+$  and open for  $\pi^-$ . The curves are from the twist-3 calculations with (line) and without (broken) sea and anti-quark contribution. predictions from the Sivers function calculation is shown as dotted lines.

on-line CNI measurements is about 60% for Run-6 (62.4 GeV) and about 45% for run 5 (200 GeV). The systematic error on the  $A_N$  measurements is estimated to be 20% including uncertainties from the beam polarization (~ 18%). The systematic error represents mainly scaling uncertainties on the values of  $A_N$ .

The rate of pions in given  $p_{\rm T}$  and  $x_F$  bins are determined in similar way as that for inclusive spectra, excepts that acceptance and efficiency corrections are not performed since all of these will cancel out in the calculation of the quantities needed for determination of  $A_N$ .

The analyzing power  $A_N$  for charged pions,  $A_N(\pi^+)$  and  $A_N(\pi^-)$  at  $\sqrt{s} = 62.4$  GeV as a function of  $x_F$  are shown in Fig. 4 using data taken at 2.3° and 3°. The  $p_{\rm T}$  range covered is from about 0.6 GeV/c at the lowest  $x_F$  up to 1.5 GeV/c at the highest  $x_F$ . The measured  $A_N$  values show strong dependence in  $x_F$  reaching large asymmetries reaching up to ~ 40% at  $x_F \sim 0.6$ . The  $A_N$  values are positive for  $\pi^+$  and negative for  $\pi^-$  and with approximate the same magnitude. The asymmetries and their  $x_F$ -dependence are qualitatively in agreement with the measurements from E704/FNAL.

The analyzing power  $A_N$  for charged pions,  $A_N(\pi^+)$  and  $A_N(\pi^-)$  at  $\sqrt{s} = 200$  GeV as a function of  $x_F$  are shown in Fig. 5 using data taken at 2.3° and 4°. The kinematic range is quite different from the 62 GeV data. The  $x_F$ -range covered is from 0.15 to 0.35, while the  $p_T$  covered by the measurements are much higher ranging for 1 to 4 GeV/c.

Figure 4 compares  $A_N(\pi)$  with a pQCD calculation in the range of  $p_T > 1 \text{ GeV}/c$  using "extended" twist-3 parton distributions [7] including the "nonderivative" contributions [19, 20]. In this framework, two calculations from the



**Fig. 5.**  $A_N$  v.s  $\mathbf{x}_F$  for pions at 200 GeV. Solid symbols are for  $\pi^+$  and open for  $\pi^-$  measured at 2.3° (left panel) and 4° (right panel). The curves are from the twist-3 calculations with (line) and without (broken) sea and anti-quark contribution. predictions from the Sivers function calculation is shown as dotted lines.

model are compared with the data: two valence densities  $(u_v, d_v)$  in the ansatz with and without sea- and anti-quark contribution in the model fit. The calculations describe the data within the uncertainties.

As the calculations shown in the figure, the dominant contribution to SSAs are from valence quarks and sea- and anti-quark contributions on SSAs are small that the current measurements are not able to quantitatively constrain the contribution. The data are also compared with Sivers mechanism which successfully describe FNAL/E704  $A_N$  data. The calculations compared with the data use valence-like Sivers functions [21,22] for u and d quarks with opposite sign. The fragmentation functions used are from the KKP parameterization [13], but the Kretzer fragmentation function [14] gives similar results. The calculations shown with dotted lines in the figure underestimate  $A_N$  for both  $p_T$  ranges, which indicates that TMD parton distributions are not sufficient to describe the SSA data at 62.4 GeV. The4 same kind of calculations are compared to the 200 GeV data in Fig.5

#### 4. Summary

In summary, unbiased invariant cross sections of identified charged particles as function of  $p_T$  were measured at high rapidity in p+p collisions at  $\sqrt{s} = 200 \text{GeV}$ and  $\sqrt{s} = 62 \text{GeV}$ . NLO pQCD calculations reproduce reasonably well the produced particle (pions and kaons) distributions. The SSA for inclusive pions were measured at forward rapidities at the same two energies. A twist-3 pQCD model of  $A_N$ describes the  $x_F$  and energy dependence reasonable well. There are theoretical challenges in describing the data at both lower energy and lower  $p_T$ .

#### Acknowledgments

We thank Werner Vogelsang, Feng Yuan and Umberto D'Alesio for providing us with their calculations shown in this contribution and in forth coming publications. This work is supported by the Division of Nuclear Physics of the Office of Science of the U.S. Department of energy under contract DE-AC02-98-CH10886. A special thank to Renaissance Technology for its generous supplemental support that allowed for the RHIC pp run in 2006.

### References

- 1. I. Arsene et al., BRAHMS Collaboration, Nucl. Phys. A.757 1 (2005)
- 2. B. Back et al., PHOBOS Collaboration, Nucl. Phys. A.757 28 (2005)
- 3. J. Adams et al., STAR Collaboration, Nucl. Phys. A.757 102 (2005)
- 4. J. Acox et al., PHENIX Collaboration, Nucl. Phys. A.757 184 (2005)
- 5. D.Sivers, *Phys.Rev.***D41** 83 (1990)
- 6. J.C.Collins, Nucl. Phys. B396 161 (1993)
- 7. J. Qiu and G. Sterman, Phys. Rev. D59 014004 (1999).
- For a review, see J. Kodaira and K. Tanaka, Prog. Theor. Phys. 101 191 (1999).
- 9. R. Debbe et al., Nucl. Instrum. Methods Phys. Res., Sect. A570, 216 (2007).
- M. Adamczyk *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A499, 437 (2003).
- 11. W. Vogelsang, private communication.
- 12. J. Pumplin et al., JHEP 0207, 012 (2002); arXiv:hep-ph/0201195.
- 13. B.A. Kniehl et al., Nucl. Phys. B597, 337 (2001).
- 14. S. Kretzer *et al*, Eur. Phys. J. **C22**, 269 (2001).
- 15. S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 241803 (2003).
- J. Adams *et al.*, Phys. Rev. Lett. **92**, 171801 (2004).
  Phys. Rev. D **67**, 054005 (2003).
- 17. I. Arsene et al., BRAHMS Collaboration. arXiv:hep-ex/0701041
- 18. C. Bourrely and J. Soffer, Eur. Phys. J. C36 371 (2004).
- 19. C. Kouvaris et al., Phys. Rev. D74 1104013 (2006).
- 20. The calculations were provided by F. Yuan.
- U. D'Alesio and F. Murgia, *Phys. Rev.* D70 074009 (2004), SPIN06 proceedings.
- 22. The calculations were provided by U. D'Alesio.